TECHNICAL DATA SHEET

FOLICAL 19

FOLIAR

PRODUCT CATEGORY

Rapidly available calcium

ELEMENTAL ANALYSIS:

Elemental Analysis:	%W/V
Calcium (Ca)	19.0

KEY BENEFITS:

- Corrects calcium deficiencies
- Strengthens cell walls in fruit and vegetables
- Effective treatment for all crops

APPLICATION RATES & TIMING:

Crop	Rate L/ha	Timing
Almond	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Apple	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Avocado	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Banana	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Brassica	5 L/ha	As required.
Broadacre	2-5 L/ha	As required.
Canola	2-5 L/ha	As required.
Carrot	5 L/ha	As required.
Cereals	2-5 L/ha	As required.
Cherry	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Citrus	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Cotton	2-5 L/ha	Pre-flower where Ca deficiency exists.
Green Bean	5 L/ha	As required.
Lettuce	5 L/ha	As required.
Legumes, Field Peas	2-5 L/ha	As required.
Lucerne	5 L/ha	As required.
Macadamia	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Mango	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Melons	5 L/ha	As required.
Onion	5 L/ha	As required.
Orchard Crops	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Potatoes	5 L/ha	As required.
Pumpkin	5 L/ha	As required.
Stone Fruit	5-7 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Strawberry	5 L/ha	As required.
Sugar Cane	5 L/ha	As required.

Continued overleaf

Sweet Potato	5 L/ha	As required.
Table Grape	5 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.
Tomato	5 L/ha	As required.
Turf	5 L/ha	As required.
Vegetables	5 L/ha	As required.
Wine Grape	5 L/ha	Pre-flower where Ca deficiency exists and every 1-2 weeks until end of cell division.

APPLICATION GUIDELINES:

MIX WELL BEFORE USE

High analysis flowable and available, low pH for improved uptake.

Can be applied by boom spray, aerial spray or fertigation.
Water rates will vary according to crop type, growth stage and canopy area.

If mixing with other products, test for compatibility with a small quantity in a jar.

TYPICAL CHEMICAL CHARACTERISTICS:

pH: 2.5
Colour: Light yellow
Form: Liquid
Odour: Odourless
Density @ 20°C: 1.35 g/cm^3
Solubility in Water: Fully miscible

PACKAGING SIZES AVAILABLE:

20 L, 200 L and 1000 L

IDEAL STORAGE CONDITIONS:

Store in cool and dry conditions (Preferably below 30°C)

© Copyright 2025. Omnia Specialities (Australia) Pty Ltd.

Disclaimer: Please note the information contained in this Technical Data Sheet (TDS) is provided for general guidance and is believed to be accurate and reliable at the time of publication. Although reasonable care and diligence have been exercised with the compilation of this TDS, and the information is given in good faith, the user assumes all risks associated with the use of this information and the use, handling, storage, and disposal of our products. In no event shall Omnia, its affiliates, or its employees be liable for any direct, indirect, incidental, special, or consequential damages arising out of the use or inability to use the information provided in this Technical Data Sheet. This document may be subject to change without notice.

FCALTDS300125