TECHNICAL DATA SHEET

OMNIBOOST-K®

FOLIAR

PROUDLY MADE IN AUSTRALIA

PRODUCT CATEGORY

Organic Based Liquid Foliar Feed

ELEMENTAL ANALYSIS:

Elemental Analysis:	%W/V
Nitrogen (N)	6.00
Phosphorous (P)	13.00
Potassium (K)	3.10
Sulphur (S)	3.30
Magnesium (Mg)	2.00
Manganese (Mn)	2.90
Iron (Fe)	0.26
Copper (Cu)	0.34
Zinc (Zn)	2.00
Boron (B)	0.35
Molybdenum (Mo)	0.04
Cobalt (Co)	0.02
Amino Acids, K-Fulvate (Fulvic Acid), Uptake Enhancers	

KEY BENEFITS:

- Boost root growth
- Improve drought resistance
- Overcome nutrient deficiencies
- Stimulate cell division
- Increase crop yields
- Improve profitability

APPLICATION RATES & TIMING:

Crop	Rate L/ha	Timing
Almond	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until hull split.
Apple	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Avocado	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Banana	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Brassica	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Broadacre	2-5 L/ha	Apply during early plant development and beginning of reproductive cycle.
Canola	2-5 L/ha	Apply during early plant development and beginning of reproductive cycle.
Carrot	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Cereals	2-5 L/ha	Apply during early plant development and beginning of reproductive cycle.
Cherry	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Citrus	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Cotton	5 L/ha	Apply from row closure and early boll development.
Green Bean	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.

Continued overleaf

Lettuce	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Legumes, Field Peas	2-5 L/ha	Apply during early plant development and beginning of reproductive cycle.
Lucerne	2-5 L/ha	Apply at 15cm growth after each cut.
Macadamia	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Mango	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Melons	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Onion	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Orchard Crops	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Potatoes	5-7 L/ha	2-3 sprays 2 weeks apart from 40mm king tuber.
Pumpkin	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Stone Fruit	5-7 L/ha	2 sprays 4 weeks apart from end of fruit cell division until ripening begins.
Strawberry	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Sugar Cane	5 L/ha	2 sprays 4 weeks apart from mid growth
Sweet Potato	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Table Grape	3-5 L/ha	2 sprays from pea size until verasion.
Tomato	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Turf	5 L/ha	Repeat every 2-4 weeks during growing season or during stress events.
Vegetables	5 L/ha	Repeat every 2-3 weeks during growing season after canopy establishment or during stress events.
Wine Grape	3-5 L/ha	2 sprays from pea size until verasion.

APPLICATION GUIDELINES:

MIX WELL BEFORE USE

For best results, apply when plants have sufficient leaf area.

Suitable for use with all forms of foliar application systems. For best results, apply when plants have sufficient leaf area.

If mixing with other products, test for compatibility with a small quantity in a jar.

TYPICAL CHEMICAL CHARACTERISTICS:

pH:	1.2
Colour:	Greenish Brown
Form:	Liquid
Odour:	Slight pungent
Density @ 20°C:	1.35 g/cm^3
Solubility in Water:	Fully miscible

PACKAGING SIZES AVAILABLE:

20 L, 200 L and 1000 L

IDEAL STORAGE CONDITIONS:

Store in cool and dry conditions (Preferably below 30°C)

® OmniBoost-K is a registered trademark of Omnia Specialities (Australia) Pty Ltd.

© Copyright 2024. Omnia Specialities (Australia) Pty Ltd.

Disclaimer: Please note the information contained in this Technical Data Sheet (TDS) is provided for general guidance and is believed to be accurate and reliable at the time of publication. Although reasonable care and diligence have been exercised with the compilation of this TDS, and the information is given in good faith, the user assumes all risks associated with the use of this information and the use, handling, storage, and disposal of our products. In no event shall Omnia, its affiliates, or its employees be liable for any direct, indirect, incidental, special, or consequential damages arising out of the use or inability to use the information provided in this Technical Data Sheet. This document may be subject to change without notice.

OBKTDS210524