omnia NUTRIOLOGY

PROUDLY MADE IN AUSTRALIA

OMNIKELP®

SOIL & FOLIAR

ACIDIC SEAWEED CONCENTRATE

1000L

NASAA INPUT FOR ORGANIC PRODUCTION Cert. No. 3456M

Scan to access full SDS

APPLICATION RATES

Crop	Rate (L/ha)	Application Foliar
Broadacre Crops	2 L/ha	Apply during early plant development and beginning of reproductive cycle.
Cotton	3 L/ha	Apply during early plant development and early boll development.
Orchard Crops	3-5 L/ha	4 sprays 4 weeks apart from early canopy development until end of fruit cell division and immediately post harvest.
Potatoes	5 L/ha	2-3 sprays two weeks apart from early emergence.
Sugar Cane	3 L/ha	2 sprays 4 weeks apart during early growth.
Vegetables	3 L/ha	2-3 sprays 2-3 weeks apart starting at establishment.
Vineyards	3 L/ha	4 sprays, 2 cm shoot growth, 10cm shoot growth and pre flowering, as required and immediately post harvest.
Crop	**Rate (L/ha)**	**Application Soil**
All Crops	5-10 L/ha	Apply from early in the crop cycle.

DIRECTIONS FOR USE

MIX WELL BEFORE USE

For best results apply OmniKelp® before fertilizer application.

Suitable for use with all forms of foliar application systems. For best results, apply when plants have sufficient leaf area.

Do not store mixed product.

Scan to access TDS/Technical Data Sheet

Elemental Analysis:	%W/V
Kelp Extracts	9.0-10.0
Phosphorous (P)	1.5
Potassium (K)	1.3
Auxins, cytokynins, jasmonates, phenolics, amino acids	

CONDITIONS OF SALE

Every care is taken during the manufacture of this product. However, as environmental conditions can vary considerably, no responsibility can be taken by the manufacturer, importer or his agents for any injury, damage, loss or other results following the use or misuse of this product.

OKELACI1000L240725

Omnia Specialities Australia Pty Ltd
(ACN 102 717 019)
Tramway Road, Morwell 3840, Victoria, Australia
Tel: +61 3 5133 9118 **E:** info@omnia.net.au
omnia.com.au

omnia NUTRIOLOGY